



Dermatology Reports

<https://www.pagepress.org/journals/index.php/dr/index>

eISSN 2036-7406



ASSOCIAZIONE DERMATOLOGI-VENEROLOGHI
OSPEDALIERI ITALIANI e della SANITÀ PUBBLICA



SIDCO

Società Italiana di Dermatologia
Chirurgica, Oncologica, Correttiva ed Estetica

Publisher's Disclaimer. E-publishing ahead of print is increasingly important for the rapid dissemination of science. **Dermatology Reports** is, therefore, E-publishing PDF files of an early version of manuscripts that undergone a regular peer review and have been accepted for publication, but have not been through the copyediting, typesetting, pagination and proofreading processes, which may lead to differences between this version and the final one. The final version of the manuscript will then appear on a regular issue of the journal. E-publishing of this PDF file has been approved by the authors.

*Please cite this article as: **Almuntrbellah Almudimeegh, Khalid Nabil Nagshabandi, Mae Barakeh, Turkey Alsehli. A rare case of Unilateral Nevoid Telangiectasia in a Saudi pediatric patient. Dermatol Rep 2024 [Epub Ahead of Print] doi: 10.4081/dr.2024.10091***

Submitted - Accepted

Note: The publisher is not responsible for the content or functionality of any supporting information supplied by the authors. Any queries should be directed to the corresponding author for the article.

All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article or claim that may be made by its manufacturer is not guaranteed or endorsed by the publisher.

A rare case of Unilateral Nevoid Telangiectasia in a Saudi pediatric patient

Almunturbellah Almudimeegh, Khalid Nabil Nagshabandi, Mae Barakeh, Turkey Alsehli

Department of Dermatology, College of Medicine, King Saud University, Riyadh, Saudi Arabia

Corresponding author: Khalid Nabil Nagshabandi, Department of Dermatology, College of Medicine, King Saud University, Riyadh, Saudi Arabia.

E-mail: khalidnagshabandi@gmail.com

Key words: Unilateral Nevoid Telangiectasia, UNT, vascular lesion, vascular anomaly.

Authors' contributions: all the authors made a substantive intellectual contribution. All the authors have read and approved the final version of the manuscript and agreed to be held accountable for all aspects of the work.

Conflict of interest: the authors declare no potential conflict of interest.

Funding: none.

Informed consent: written informed consent was obtained from the patient for publication of the details of their medical case and any accompanying images.

Availability of data and materials: all data generated or analyzed during this study are included in this article. Further enquiries can be directed to the corresponding author.

Abstract

Unilateral Nevoid Telangiectasia (UNT) is a rare congenital or acquired benign cutaneous vascular anomaly, first identified by Blaschko in 1899. It is characterized by superficial telangiectasias with a reticular or linear pattern on one side of the body, often following a dermatomal distribution. We present the first case documented in Saudi Arabia of UNT in a 12-year-old girl.

Introduction

Unilateral Nevoid Telangiectasia (UNT), is a rare congenital or acquired benign cutaneous vascular anomaly. First identified by Blaschko in 1899, it is characterized by grouped superficial telangiectasias with a reticular or linear net-like pattern confined to one side of the body (unilateral), that blanch with diascopy and frequently follow a dermatomal pattern.¹ The condition can present at any age but is most often identified during adolescence or early adulthood. UNT tends to occur more frequently in females than in males, mostly appearing on the upper trunk and extremities.^{1,2} The precise cause of UNT is not fully understood, though it is believed to involve both genetic and hormonal factors. Hormonal changes during puberty, pregnancy, or the use of oral contraceptives have been linked to the condition, suggesting a role for estrogen sensitivity.³⁻⁵ Additionally, UNT has been associated with systemic conditions such as liver disease and hyperthyroidism in some patients.^{2, 6-9}

Clinically, UNT presents as clusters of fine, red to purple linear or spider-like vessels that blanch under pressure. These telangiectasias are typically asymptomatic, though some patients may have mild discomfort or cosmetic concerns.¹ Diagnosis is primarily clinical, based on the characteristic appearance and distribution of the lesions. Dermoscopy can assist in the evaluation by revealing a network of non-melanocytic red tortuous dilated capillaries.¹⁰ Despite its benign nature, UNT can sometimes be mistaken for other vascular or dermatologic conditions, highlighting the importance of awareness and accurate diagnosis among dermatologists. Differential diagnoses include port-wine stains, spider angiomas, and other causes of unilateral vascular lesions. In this report, we present the first documented case of a Saudi pediatric patient diagnosed with unilateral nevoid telangiectasia, a condition previously unreported in Saudi Arabia.

Case Report

We report the case of a healthy 12-year-old girl, who presented to the Dermatology outpatient clinic complaining of multiple red-violet cutaneous lesions on her right forearm. The lesions began at birth and have grown with her age. They were asymptomatic, with no previous bleeding, ulceration, infections, or change in size. In addition, her personal and family medical history was insignificant. The only concern is their appearance.

Clinical examination revealed multiple ill-defined violaceous blanchable macules that disappear with pressure (**Figure 1**), involving the lower lateral segment of the right dorsal forearm with some proximal extension (**Figure 2**). The rest of the examination was normal, with no mucosal lesions. We proposed a diagnosis of unilateral nevoid telangiectasia (UNT). Based on these findings, the patient was given follow-up visits to receive Pulsed-dye laser sessions.

Discussion

Unilateral nevoid telangiectasia is a rare and uncommon cutaneous condition with an unknown etiology. It is clinically characterized by multiple blanchable red-violet telangiectasias mainly located unilaterally in a C3-T1 dermatomal pattern.^{1,11} The pathophysiology of unilateral nevoid UNT remains incompletely understood, but several contributing factors have been identified, suggesting a multifactorial etiology.

Genetic predisposition plays a significant role in the development of UNT where it often appears to follow a dermatomal distribution, which aligns with the patterns of skin development and innervation. This dermatomal presentation hints at a possible mosaicism, where genetic mutations during embryogenesis could lead to localized vascular anomalies.¹² Cases of UNT have also been associated with other rare conditions, such as ipsilateral melorheostosis and vascular twin spotting,

further indicating a possible role of somatic mosaicism.^{13,14} These genetic changes might result in the abnormal proliferation or dilation of superficial capillaries and venules, manifesting as the characteristic telangiectasias seen in UNT.¹² Hormonal influences are crucial in the pathogenesis of UNT, which is more frequently diagnosed in females and often coincides with hormonal fluctuations such as puberty, pregnancy, and oral contraceptive use. Estrogen, in particular, increases the synthesis of vascular endothelial growth factor (VEGF) and other angiogenic factors, promoting blood vessel formation and dilation.⁹

Systemic conditions may also contribute to UNT development. Associations have been observed with systemic diseases like liver cirrhosis and malignancy, chronic alcoholism, hepatitis B & C virus infection, and hyperthyroidism.^{2, 6-9} These conditions may worsen vascular abnormalities by altering blood flow dynamics, affecting hormone metabolism, or having a direct impact on vascular integrity. Liver diseases can elevate circulating estrogen levels due to impaired metabolism hence increasing the levels of VEGF, which potentially triggers telangiectatic changes in susceptible individuals.⁹ Moreover, UNT has been observed in conjunction with neurological disorders in a subset of patients, suggesting a possible neurocutaneous link. A study involving eight patients with UNT reported hypoesthesia over the skin lesions in all patients. Additionally, cranial magnetic resonance imaging revealed subcortical hamartomatous lesions in one patient and demyelinated plaques in another. Neurological symptoms in these patients included mild mental retardation, dysarthric speech, hemiparesis, and dysmetria, indicating significant neurological involvement. In one case, UNT was accompanied by findings suggestive of multiple sclerosis.¹⁵ A rare instance reported the occurrence of UNT in a healthy male following chemotherapy for Hodgkin's disease. This case suggests a potential new clinical association between chemotherapy and UNT.¹⁶ In another instance reported, UNT occurred in a healthy male without any evident

predisposing factors or lab abnormalities.¹⁷ Our case revealed the occurrence of congenital UNT in a young girl, as the lesions have been present since birth. Treatment of UNT is often sought for cosmetic reasons, and several options are available. Pulsed Dye Laser (PDL) is the primary and effective therapeutic option that showed significant improvement in the appearance of telangiectasias with minimal adverse effects.^{18,19}

Conclusions

UNT is a complex and multifactorial vascular dermatosis. Genetic mosaicism likely sets the stage for this localized condition, which is then influenced by hormonal factors and potentially exacerbated by systemic conditions.

References

1. Wenson SF, Jan F, Sepehr A. Unilateral nevoid telangiectasia syndrome: a case report and review of the literature. *Dermatol Online J.* 2011;17:2.
2. Sukanya G, Sane RR, Ravi VN, Nehete SS. Unilateral Nevoid Telangiectasia: An Overlooked Entity. *Indian J Dermatol.* 2022;67:423-4.
3. Herzum A, Micalizzi C, Parodi A. Unilateral Nevoid Telangiectasia After Pregnancy. *Dermatol Pract Concept.* 2021;11:e2021119.
4. Nohara T, Natsuga K, Yasuoka A, Ujiie H. Unilateral Naevoid Telangiectasia Associated with Oral Contraceptive. *Acta Derm Venereol.* 2021;101:adv00595.
5. Coe JS, George SMC. Image Gallery: Unmasking of unilateral naevoid telangiectasia during puberty. *Br J Dermatol.* 2018;178:e331.
6. Kim HJ, Kim KO, Kim YS, et al. Acquired Unilateral Nevoid Telangiectasia Accompanied by Liver Metastasis of Colon Cancer. *Ann Dermatol.* 2016;28:404-5.
7. Karakaş M, Durdu M, Sönmezoğlu S, et al. Unilateral nevoid telangiectasia. *J Dermatol.* 2004;31:109-12.
8. Kavak A, Kutluay L. Unilateral nevoid telangiectasia and hyperthyroidism: a new association or coincidence? *J Dermatol.* 2004;31:411-4.
9. Smith JA, Kamangar F, Prakash N, et al. Unilateral nevoid telangiectasia syndrome (UNTS) associated with chronic hepatitis C virus and positive immunoreactivity for VEGF. *Dermatol Online J.* 2014;20:13030/qt8g9268sf.
10. Villela-Segura U. Dermoscopy as an Important Tool for Differentiating Unilateral Nevoid Telangiectasia and Angioma Serpiginosum. *Dermatol Pract Concept.* 2019;9:306-7.

11. Karabudak O, Dogan B, Taskapan O, Harmanyeri Y. Acquired unilateral nevoid telangiectasia syndrome. *J Dermatol.* 2006;33:825-6.
12. van Steensel MA. Neurocutaneous Manifestations of Genetic Mosaicism. *J Pediatr Genet.* 2015;4:144-53.
13. Kim J, Cho SB, Cho S, Bang D. Unilateral nevoid telangiectasia associated with ipsilateral melorheostosis. *Ann Dermatol.* 2012;24:206-8.
14. Tan C, Zhu WY. Unilateral nevoid telangiectasia superimposed on the Bier spots: another demonstration of vascular twin spotting. *J Dtsch Dermatol Ges.* 2011;9:389-90.
15. Akman-Karakaş A, Kandemir H, Senol U, et al. Unilateral nevoid telangiectasia accompanied by neurological disorders. *J Eur Acad Dermatol Venereol.* 2011;25:1356-9.
16. Rodríguez-Martín M, Sáez M, Carnerero A, et al. Unilateral naevoid telangiectasia in a young man after chemotherapy: a simple coincidence or a new clinical association? *J Eur Acad Dermatol Venereol.* 2006;20:1001-2.
17. Jordão JM, Haendchen LC, Berestinas TC, Faucz LR. Acquired unilateral nevoid telangiectasia in healthy men. *An Bras Dermatol.* 2010;85:912-4.
18. Sharma VK, Khandpur S. Unilateral nevoid telangiectasia--response to pulsed dye laser. *Int J Dermatol.* 2006;45:960-4.
19. An XJ, Yang J, Gao Y, et al. Unilateral nevoid telangiectasia treated with pulsed dye laser: Use of dermoscopy to monitor the response. *Chin Med J (Engl).* 2019;132:2125-6.



Figure 1. Multiple ill-defined violaceous macules.



Figure 2. Unilateral Nevoid Telangiectasia (UNT) involves the lower lateral segment of the forearm and extends proximally.