

## Skin reaction to COVID-19 vaccine: A report of 4 cases

Mariabeatrice Bertolani,<sup>1</sup>

Maria Beatrice de Felici del Giudice,<sup>1</sup>

Erminia Ridolo,<sup>2</sup> Francesco Pucciarini,<sup>2</sup>

Claudio Feliciani,<sup>1</sup> Francesca Satolli<sup>1</sup>

<sup>1</sup>Section of Dermatology, Department of Medicine and Surgery, and <sup>2</sup>Medicine and Surgery Department, Allergy and Clinical Immunology, University of Parma, Italy

### Abstract

Since the beginning of the covid-vaccine campaign, a lot of local and systemic dermatologic reactions happening after the administration of Coronavirus disease 2019 (COVID-19) vaccines have been described, even if their exact biological mechanism is still debated. In this paper we report 4 cases of cutaneous manifestations arose within ten days after the first dose of messenger RNA (mRNA)-based COVID-19 vaccination: one case of giant urticaria, one case of head and neck redness and two cases of Erythema Multiforme (EM). In our experience these reactions were mild, transient and all of them resolved, not recurring after the second dose, so these manifestations shouldn't be considered as an absolute contraindication to the second dose of vaccine, that to date is fundamental.

### Introduction

As of today, the Italian Drugs Agency (AIFA) has approved four different SARS-CoV-2 vaccines: Pfizer/BioNTech (mRNABNT162b2, Comirnaty) and Moderna (mRNA-1273) are messenger RNA (mRNA) vaccines, while AstraZeneca (AZD1222/ChAdOx1, Vaxzevria) and Janssen (Ad26.COV2.S) are two viral vector vaccines.<sup>1</sup> It's by now well-known that Coronavirus disease 2019 (COVID-19) has numerous and various cutaneous manifestations: since the beginning of the pandemic one and a half year ago a large range of skin manifestations has been associated with the infection, such as chilblain-like, urticarial, vesicular, maculopapular, livedoid and vasculitic lesions. Nevertheless, the exact pathophysiology of these manifestations is still not totally clear; various hypotheses have been proposed, including viral hypersensitivity reactions, virus induced coagulopathy, thrombotic microangiopathy,

overexpression of type I interferons and direct viral damage. It stands to reason that skin reactions can occur even after vaccination especially with mRNA vaccines encoding the spike protein of SARS-CoV-2.<sup>2</sup>

We report 4 cases of cutaneous manifestations which appeared within ten days after the first dose of an mRNA COVID-19 vaccination (Table 1).

### Case report #1

A 58-year-old man was administered the first dose of the mRNA COVID-19 vaccine Pfizer/BioNTech. In the four days following the vaccination, the patient developed severe itching with fleeting urticarial lesions lasting <24 hours and migrant in appearance with polycyclic contours associated with hand swelling. No angioedema nor respiratory symptoms were present. Laboratory investigations detected raised PCR (18.4 mg/dL) and white cell count (10.410/microL). The diagnosis was consistent with giant urticaria (Figures 1a,b). The patient was treated with 0.5 mg/kg methylprednisolone daily plus oral antihistamine which resulted in the complete resolution of skin lesions within 10 days.

### Case report #2

A 59-year-old man presented to the emergency department two days after the first dose of the Moderna Covid-19 vaccine for precordial pain associated with tachycardia, paresthesia and light-headedness. In a clinical examination he presented with a persistent head and neck redness (Figures 1c,d) associated with skin-colored, well-defined macules with a peripheral keratotic ridge, which were widespread on arms and legs bilaterally. His past medical history included psoriasis, which had always been treated with a topical combination of calcipotriol/betamethasone, and anxiety for which he took Alprazolam daily. Our differential diagnoses comprised head and neck dermatitis, airborne dermatitis, photoinduced dermatitis but he had no personal nor family history of atopy nor autoimmune disorders. He was treated with oral Cetirizine 10mg/die and slowly improved.

### Case reports #3 and #4

A 22-year-old girl presented with targetoid lesions symmetrically distributed

Correspondence: Mariabeatrice Bertolani, Section of Dermatology, Department of Clinical and Experimental Medicine, University of Parma, Via Gramsci 14, 43126 Parma, Italy.  
Tel.: +39-0521-702711, fax: +39-0521-702959  
E-mail: mariabeatrice.bertolani@gmail.com

Key words: COVID-19; Vaccination; Skin reaction.

Contributions: MB and MBFG contributed equally.

Conflict of interest: The authors declare no potential conflict of interest.

Funding: None.

Availability of data and material: Data and materials are available by the authors.

Please cite this article as: Bertolani M, de Felici del Giudice MB, Ridolo E, et al. Skin reaction to COVID-19 vaccine: a report of 4 cases. *Dermatol Rep* 2022;14:9376.

Received for publication: 9 September 2021.

Revision received: 18 September 2021.

Accepted for publication: 12 October 2021.

This work is licensed under a Creative Commons Attribution-NonCommercial 4.0 International License (CC BY-NC 4.0).

©Copyright: the Author(s), 2022

Licensee PAGEPress, Italy

*Dermatology Reports* 2022; 14:9376

doi:10.4081/dr.2022.9376

*Publisher's note: All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article or claim that may be made by its manufacturer is not guaranteed or endorsed by the publisher.*

on the palms and soles associated with an initial involvement of the thighs (Figure 2a,b). The patient was asymptomatic, blood test analyses were within normal values and IgM anti-HSV1, anti-HSV2, anti-Mycoplasma pneumoniae, anti-CMV, anti-EBV and anti-parvovirus B19 were all negative. Ten days previously, she had received the first dose of the Moderna Covid-19 vaccine.

Similarly, a 32-year-old man presented with pruritic concentric annular lesions bilaterally spread on the soles, palms and back of the hands, which had arisen eight days after the first dose of the Moderna Covid-19 vaccine (Figure 2c,d). Besides itching he did not complain of any systemic

symptoms. In both patients oral and genital mucosa were spared and the clinical features suggested Erythema Multiforme (EM). The former patient was successfully treated with 0.5 mg/kg methylprednisolone daily plus oral antihistamine while the latter completely resolved within one week with oral Cetirizine 10mg/die.

All patients received the second booster of vaccine without any systemic symptoms nor skin problems.

### Discussion

A large variety of both local and systemic skin reactions have recently been reported after Covid-19 vaccination, especially occurring after the mRNA types.<sup>3</sup>

The exact biological mechanism of these reaction is still debated, and, to date, it has not been possible to establish if the main cause is a hypersensitivity to the spike protein or to other components of the vaccine.<sup>2,4</sup> However, distinguishing between immediate and delayed hypersensitive reactions is essential. In particular, when manifestations such as urticaria and angioedema occur within 4 hours from the injection, a possible contraindication for the second dose can exist, especially if the skin manifestations are associated with respiratory and/or cardiovascular symptoms.<sup>5</sup> As in case 1, the first dose-related urticaria developed in a few days, suggesting that it may not be induced by a hypersensitivity reaction to the vaccine but rather related to the host's immune response. Indeed, the patient did not exhibit any symptoms after his second dose.

Regarding EM, it is typically associated with microbial infections, especially Herpesviridae viruses or Mycoplasma pneumonia, but in a minority of cases it can be related to drugs or vaccines, mainly in the pediatric population.<sup>6</sup> The two most frequently associated vaccines are the

diphtheria-tetanus-pertussis and recombinant hepatitis B<sup>7</sup> but recently an association with Covid-19 vaccines has been recognized.<sup>8</sup>

Some authors hypothesize that although the vaccine does not cause de novo immune

mediated adverse reactions, it may trigger a pre-existing underlying dysregulated pathway through the immunologic response, inducing immune activation of both B and T lymphocytes.<sup>6,9,10</sup>



Figure 1. Case reports #1 and #2.

Table 1. Summary of the 4 cases.

	Case 1	Case 2	Case 3	Case 4
Vaccine	Pfizer/BioNTech	Moderna	Moderna	Moderna
Timing of onset	4 days	2 days	10 days	8 days
Kind of reaction	Giant urticaria	Head and neck redness	EM	EM
Age	58 years old	59 years old	22 years old	32 years old
Gender	Male	Male	Female	Male
Comorbidities	None	Psoriasis, anxiety	None	None
Duration of reaction	10 days	10 days	7 days	5 days
Treatment	0.5 mg/kg methylprednisolone daily plus oral antihistamine	Oral Cetirizine 10 mg/die	0.5 mg/kg methylprednisolone daily plus oral antihistamine	Oral Cetirizine 10 mg/die

## Conclusions

In conclusion, although we cannot demonstrate a true causal link, all the reported cases show a temporal relation between skin reactions and vaccine administrations. Our cases also show that the majority of skin reactions are mild, transient and resolve without recurring after the

second dose, as observed by other authors.<sup>11</sup> Consequently these manifestations should not be considered as an absolute contraindication to a second dose of vaccine that to date is fundamental.

Meticulous reporting is necessary and further studies based on larger cohorts are needed to better understand the pathophysiology of cutaneous involvement.

## References

1. <https://www.aifa.gov.it/web/guest/vaccini-covid-19>
2. Fernandez-Nieto D, Hammerle J, Fernandez-Escribano M, et al. Skin manifestations of the BNT162b2 mRNA COVID-19 vaccine in healthcare workers. 'COVID-arm': a clinical and histological characterization. *J Eur Acad Dermatol Venereol* 2021;35:e425-7.
3. McMahon DE, Amerson E, Rosenbach M, et al. Cutaneous reactions reported after Moderna and Pfizer COVID-19 vaccination: a registry-based study of 414 cases. *J Am Acad Dermatol* 2021;7:S0190-9622.
4. Farinazzo E, Ponis G, Zelin E, et al. Cutaneous adverse reactions after mRNA COVID-19 vaccine: early reports from Northeast Italy. *J Eur Acad Dermatol Venereol* 2021;22:10.
5. Interim Clinical Considerations for Use of mRNA COVID-19 Vaccines Currently Authorized in the United States. Centers for Disease Control and Prevention. Accessed February 15, 2021. Available at: <https://www.cdc.gov/vaccines/covid-19/info-byproduct/clinical-considerations.html>
6. Lerch M, Mainetti C, Terziroli Beretta-Piccoli B, Harr T. Current perspectives on erythema multiforme. *Clin Rev Allergy Immunol* 2018;54:177-84.
7. Zoghaib S, Kechichian E, Souaid K, et al. Triggers, clinical manifestations, and management of pediatric erythema multiforme: A systematic review. *J Am Acad Dermatol* 2019;81:813-22.
8. Ayatollahi A, Hosseini H, Firooz R, Firooz A. COVID-19 vaccines: What dermatologists should know? *Dermatol Ther* 2021;7:e15056.
9. Talotta R. Do COVID-19 RNA-based vaccines put at risk of immune-mediated diseases? In reply to "potential antigenic cross-reactivity between SARS-CoV-2 and human tissue with a possible link to an increase in autoimmune diseases". *Clin Immunol* 2021;24:108665.
10. Akinosoglou K, Tzivaki I, Marangos M. Covid-19 vaccine and autoimmunity: Awakening the sleeping dragon. *Clin Immunol* 2021;3:108721.
11. Corbeddu M, Diociaiuti A, Vinci MR, et al. Transient cutaneous manifestations after administration of Pfizer-BioNTech COVID-19 Vaccine: an Italian single-centre case series. *J Eur Acad Dermatol Venereol* 2021;35:e483-e485



Figure 2. Case reports #3 and #4.