



Dermatology Reports

<https://www.pagepress.org/journals/index.php/dr/index>

eISSN 2036-7406



Publisher's Disclaimer. E-publishing ahead of print is increasingly important for the rapid dissemination of science. **Dermatology Reports** is, therefore, E-publishing PDF files of an early version of manuscripts that undergone a regular peer review and have been accepted for publication, but have not been through the copyediting, typesetting, pagination and proofreading processes, which may lead to differences between this version and the final one. The final version of the manuscript will then appear on a regular issue of the journal. E-publishing of this PDF file has been approved by the authors.

Please cite this article as: Le Lay F, Deparis M, Freitag S, Domp martin A. Rubella virus vaccine-induced granulomas: a case in children with ataxia-telangiectasia. Dermatol Rep 2024 [Epub Ahead of Print] doi: 10.4081/dr.2024.9864

 © the Author(s), 2024
Licensee PAGEPress, Italy

Submitted: 07/10/2023 – Accepted 12/11/2023

Note: The publisher is not responsible for the content or functionality of any supporting information supplied by the authors. Any queries should be directed to the corresponding author for the article.

All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article or claim that may be made by its manufacturer is not guaranteed or endorsed by the publisher.

Rubella virus vaccine-induced granulomas: a case in children with ataxia-telangiectasia

Florine Le Lay,¹ Marianna Deparis,² Sylvie Fraitag,³ Anne Domp martin¹

¹Dermatology Department, Caen University Hospital, Caen; ²Pediatrics Department, Caen University Hospital, Caen; ³Pathology Department, Necker-Enfants Malades Hospital, Paris, France

Correspondence: Florine Le Lay Dermatology Department, CHU Caen, avenue Côte-de-Nacre, 14000 Caen, France.

Tel.: +33.231272510.

E-mail: lelay-f@chu-caen.fr

Key words: ataxia telangiectasia; immunodeficiency; rubella; granuloma; vaccine.

Contributions: the authors contributed equally.

Conflict of interest: the authors declare no potential conflict of interest.

Consent for publication: the parental consent for the publication of their child's photos has been obtained, although we are unable to identify her.

Abstract

Ataxia telangiectasia (AT) is a rare autosomal recessive primary immunodeficiency disorder (PID) resulting from a mutation in the ATM gene, which is involved in DNA repair. We describe the case of a young girl with cutaneous granulomas that developed after childhood vaccinations. Immunohistochemistry revealed granulomas induced by the rubella virus vaccine. This finding raises the question of live rubella vaccine strains safety in immunocompromised children.

Introduction

Ataxia telangiectasia (AT) is a complex disorder with multiple phenotypic features and variability in the severity of symptoms between affected patients.¹

The development of cutaneous granulomas in AT is a known phenomenon, less common than in other primary immunodeficiency disorders (PIDs) such as DICV (common variable immunodeficiency disorder) and Nijmegen breakage syndrome, and is usually considered to be a response to a microbial trigger or a consequence of immune dysregulation.²

Case report

A six-year-old girl with AT was referred to our hospital with a 4-year history of well-demarcated and infiltrated erythematous-squamous plaque measuring 5x3 cm on the anterior aspect of the left leg and a 1-year history of a similar plaque on the left forearm (**Figures 1-2**). The lesions were ulcerated and non-tender with a gradual extension. Physical examination revealed hepatosplenomegaly.

Biology showed low levels of LB CD19+, LT CD4+ and LT CD8+ with hyper IgM immunophenotype (deficiency of IgA and IgG and normal IgM), despite immunoglobulin cures started one year ago. No cytopenia was associated.

Skin biopsy excluded a neoplastic lesion and confirmed the diagnosis of non-sarcoid granuloma. The specimen showed granulomatous dermal inflammation with necrotizing epithelioid and giant cells. No infectious agent was identified by conventional methods. Abdominal ultrasound and whole-body magnetic resonance imaging showed multiple supra- and infra-diaphragmatic adenomegaly, pulmonary nodules, and hepatosplenomegaly with micronodules. The patient's parents objected to sampling of deeper lesions to confirm extracutaneous granuloma. A myelogram was performed, and blood disorders were excluded.

The child has received all required live vaccines prior to the diagnosis of PID. Immunohistochemistry (IHC) of the skin biopsy, using a monoclonal anti-rubella capsid antibody (Abcam 34749), demonstrated the presence of rubella virus capsid within the granulomas (**Figure 3**). The

diagnosis of rubella vaccine-induced cutaneous granulomas was made. The skin lesions did not respond to topical corticosteroids.

Discussion

Cutaneous granulomas in PIDs represent a recognized complication, yet their incidence in individuals with AT is relatively low, affecting less than 10% of patients.² As the search for responsible microorganisms or antigens is frequently unsuccessful, granulomas are often considered as immune dysregulation and a correlation with a hyperIgM immunoglobulin phenotype has been reported.^{3,4} Recent investigations have identified rubella as a primary virus strongly linked to cutaneous granulomas in immunocompromised children, substantiated by the absence of viral detection in biopsies from healthy skin.⁵ A case report of rubella granuloma (RG) due to wild type in AT illustrates the fact that it is not vaccine strain specific.⁶ Skin RG can appear weeks to decades after infection or vaccination, typically before the age of two, and progresses slowly over years.^{5,6,7} It is characterised by well-defined erythematous-squamous papular lesions with atrophic centres, usually localised to the trauma-prone area, which can cause significant morbidity with possible local tissue destruction.⁴ Extracutaneous lesions have been described in two studies of patients with PID, showing that the spleen, liver, bones and lungs are the most commonly affected sites.^{4,5} These findings align with our case, suggesting a potential visceral process underlying the splenic and pulmonary nodules observed in our patient.

The presence of rubella vaccine capsid proteins in granuloma M2 macrophages suggests that macrophage tropism is a trigger for granuloma formation, such as aberrant CD8+ cytotoxic T cell response, profound T-cell deficiency, and abnormal cell repair DNA.⁸

To date, no treatment was known to be highly effective to cure RG. In patient with immunodeficiency, immunoglobulin therapy has no benefit. Nitazoxanide, while exhibiting antiviral activity against rubella virus in vitro, demonstrated limited efficacy when administered to patients.⁹ Regression of RG was only seen after hematopoietic stem cell transplantation partly explained by the elimination of rubella-carrying neutrophils and monocytes.¹⁰

The diagnosis of RG is based on characteristic clinical features and skin biopsy with histopathology and IHC or PCR. In immunocompromised children, IHC or PCR should be performed on all idiopathic cutaneous granulomas to elucidate the aetiology and frequency of rubella virus involvement. Possibly, the rubella virus vaccine strain and wild type may be the etiology of cutaneous granulomatous dermatitis previously diagnosed as idiopathic.

In most cases, the diagnosis of AT is made after the period of childhood vaccinations. Vaccine-induced cutaneous granulomas have not been seen with live attenuated measles, mumps or

varicella vaccines.⁷ Although rubella vaccination is recommended despite this potential late side effect, the occurrence of RG raises concerns about its safety.²

List of abbreviations

AT : Ataxia Telangiectasia

IHC : Immunohistochemistry

RG : Rubella granuloma

PID : Primary immunodeficiency

References

1. van Os NJH, Jansen AFM, van Deuren M, Haraldsson A, van Driel NTM, Etzioni A, et al. Ataxia-telangiectasia: Immunodeficiency and survival. *Clin Immunol*. 2017 May;178:45–55.
2. Rothblum-Oviatt C, Wright J, Lefton-Greif MA, McGrath-Morrow SA, Crawford TO, Lederman HM. Ataxia telangiectasia: a review. *Orphanet J Rare Dis*. 2016 Dec;11(1):159.
3. Chiam LYT, Verhagen MMM, Haraldsson A, Wulffraat N, Driessen GJ, Netea MG, et al. Cutaneous Granulomas in Ataxia Telangiectasia and Other Primary Immunodeficiencies: Reflection of Inappropriate Immune Regulation? *Dermatology*. 2011;223(1):13–9.
4. Buchbinder D, Hauck F, Albert MH, Rack A, Bakhtiar S, Shcherbina A, et al. Rubella Virus-Associated Cutaneous Granulomatous Disease: a Unique Complication in Immune-Deficient Patients, Not Limited to DNA Repair Disorders. *J Clin Immunol*. 2019 Jan;39(1):81–9.
5. Neven B, Pérot P, Bruneau J, Pasquet M, Ramirez M, Diana JS, et al. Cutaneous and Visceral Chronic Granulomatous Disease Triggered by a Rubella Virus Vaccine Strain in Children With Primary Immunodeficiencies: Table 1. *Clin Infect Dis*. 2017 Jan 1;64(1):83–6.
6. Browne R, Cliffe L, Ip W, Brown K, McDermott E. A case of wild-type rubella-associated cutaneous granuloma in ataxia telangiectasia. *Pediatr Dermatol*. 2022 Jul;39(4):619–21.
7. Groß M, Speckmann C, May A, Gajardo-Carrasco T, Wustrau K, Maier SL, et al. Rubella vaccine-induced granulomas are a novel phenotype with incomplete penetrance of genetic defects in cytotoxicity. *J Allergy Clin Immunol*. 2022 Jan;149(1):388-399.e4.
8. Perelygina L, Plotkin S, Russo P, Hautala T, Bonilla F, Ochs HD, et al. Rubella persistence in epidermal keratinocytes and granuloma M2 macrophages in patients with primary immunodeficiencies. *J Allergy Clin Immunol*. 2016 Nov;138(5):1436-1439.e11.
9. Perelygina L, Buchbinder D, Dorsey MJ, Eloit M, Hauck F, Hautala T, et al. Outcomes for Nitazoxanide Treatment in a Case Series of Patients with Primary Immunodeficiencies and Rubella Virus-Associated Granuloma. *J Clin Immunol*. 2019 Jan;39(1):112–7.

10. Baumann U, Schulte JH, Groß JP, Beier R, Ludwig M, Wahn V, et al. Case Report: Rubella Virus-Induced Cutaneous Granulomas in Two Pediatric Patients With DNA Double Strand Breakage Repair Disorders – Outcome After Hematopoietic Stem Cell Transplantation. *Front Immunol.* 2022 Jun 2;13:886540.

Figures



Figure 1. Macroscopic aspect of left leg cutaneous granuloma.



Figure 2. Macroscopic aspect of left forearm cutaneous granuloma.




Figure 3. Immunohistochemical study with anti-Rubella antibody showing numerous positive foci within the granulomas (magnification x5).