

SUPPLEMENTARY MATERIALS

Platelet-rich plasma for the treatment of scleroderma-associated ulcers: a single-center experience and literature review

Alessandra Grazia Condorelli,^{1*} Alessia Paganelli,^{1*} Chiara Marraccini,² Elena Ficarelli,¹ Alfonso Motolese,³ Luca Contu,¹ Alberico Motolese¹

¹Dermatology Unit, Azienda USL-IRCCS di Reggio Emilia, Arcispedale Santa Maria Nuova, Reggio Emilia; ²Transfusion Medicine Unit, AUSL-IRCCS di Reggio Emilia; ³Section of Dermatology, Department of Clinical and Experimental Medicine, University of Messina, Italy

*These authors contributed equally.

Correspondence: Alessandra Grazia Condorelli, Arcispedale Santa Maria Nuova of Reggio Emilia, Viale Risorgimento 80, 42123, Reggio Emilia, Italy.

Tel.: +39.3335359202.

E-mail: alecondor91@outlook.it

Key words: PRP, scleroderma ulcers, homologous PRP, systemic sclerosis.

Conflict of interest: the authors declare no potential conflict of interest.

Funding: none.

Availability of data and material: data and materials are available by the authors.

Informed consent: the patients involved in this study gave their consent for the publication.

Table S1. Scleroderma skin ulcers features (size and site), before treatment and after the end of treatment. Number of weeks necessary to healing process.

| Skin Ulcers | Size, cm ² | | Site | Healing time (weeks) |
|----------------|-----------------------|-----------|--------|-------------------------|
| | Before PRP | After PRP | | |
| 1 | 2,94 | 0,19 | Finger | 3 |
| 2 | 28,26 | 3,14 | Leg | 4 |
| 3 | 9,42 | 1,57 | Leg | 5 |
| 4 | 5,88 | 0,19 | Leg | 5 |
| 5 | 0,78 | 0,19 | Heel | 6 |
| 6 | 15,7 | 1,17 | Leg | 8 |
| 7 | 1,57 | 1,33 | Finger | 3 |
| 8 | 2,35 | 0,23 | Finger | 4 |
| 9 | 2,94 | 0,78 | Finger | 5 |
| 10 | 5,88 | 1,57 | Finger | 3 |
| Mean | 7,57 | 1,04 | | 4.7 |

PRP, platelet-rich plasma.