



Dermatology Reports

<https://www.pagepress.org/journals/index.php/dr/index>

eISSN 2036-7406



Publisher's Disclaimer. E-publishing ahead of print is increasingly important for the rapid dissemination of science. **Dermatology Reports** is, therefore, E-publishing PDF files of an early version of manuscripts that undergone a regular peer review and have been accepted for publication, but have not been through the copyediting, typesetting, pagination and proofreading processes, which may lead to differences between this version and the final one. The final version of the manuscript will then appear on a regular issue of the journal. E-publishing of this PDF file has been approved by the authors.

Please cite this article as: Maione V, Miccio L, Rovaris S, et al. Eosinophilic fasciitis and breast cancer: a case report highlighting recurrence signals. Dermatol Rep 2023 [Epub Ahead of Print] doi: 10.4081/dr.2024.9928

 © the Author(s), 2024
Licensee PAGEPress, Italy

Submitted: 05/01/2024 – Accepted 12/01/2024

Note: The publisher is not responsible for the content or functionality of any supporting information supplied by the authors. Any queries should be directed to the corresponding author for the article.

All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article or claim that may be made by its manufacturer is not guaranteed or endorsed by the publisher.

Eosinophilic fasciitis and breast cancer: a case report highlighting recurrence signals

Vincenzo Maione,¹ Laura Miccio,² Sara Rovaris,¹ Davide Zardo,² Stefano Bighetti,¹ Luca Bettolini,¹ Luigi Naldi²

¹Department of Dermatology, University of Brescia, ASST Spedali Civili di Brescia, Brescia;

²Department of Dermatology, San Bortolo Hospital, Vicenza, Italy

Correspondence: Luca Bettolini, Department of Dermatology, Spedali Civili, University of Brescia, Brescia, Italy

Tel.: +39.030 3995301.

Fax: +39.0303995015.

E-mail: lbettolini@gmail.com

Key word: eosinophilic fasciitis; paraneoplastic syndrome; breast cancer recurrence.

Conflict of interest: the authors declare no potential conflict of interest.

Funding: none.

Ethics approval and consent to participate: no ethical committee approval was required for this case report by the Department, because this article does not contain any studies with human participants or animals. Informed consent was obtained from the patient included in this study.

Consent for publication: the patient in this manuscript has given written informed consent to publication of her case details.

Abstract

Eosinophilic Fasciitis (EF) poses a diagnostic challenge owing to its uncommon occurrence and aetiology that remains poorly understood. We report a case of a 79-year-old woman with a prolonged history of breast cancer exhibiting EF with concomitant pelvic metastases. This case underscores EF's potential as an indicator signaling a potential resurgence of cancer, highlighting its significance as a red flag of disease recurrence.

Introduction

Eosinophilic Fasciitis, also known as Shulman disease, stands as a rare sclerotic affliction affecting muscular fasciae. Despite its clinical significance, the etiology remains elusive, and the pathogenesis remains a subject of limited comprehension. This enigmatic disorder has been observed in correlation with an array of conditions, encompassing hematologic disorders, solid neoplasms, autoimmune diseases, drugs, and infections¹.

Case report

We present a case of a 79-year-old woman with a bothersome, itchy skin rash and distinctive symptoms in her left leg: erythema, edema, and a "peau d'orange" texture. Despite no signs of infection (Fig. 1a), she had a history of breast cancer spanning 19 years. Her treatment initially involved an aromatase inhibitor, then capecitabine, which was stopped due to hand-foot syndrome. Following this, she noticed leg edema and subsequent sclerosis. Over three months, her eosinophil count rose (from 0.51/mm³ to 1.3/mm³), and platelet count at 438000/mm³. An echocolor Doppler scan ruled out thrombosis.

A skin biopsy of the left leg revealed hyperkeratotic epidermis, dermal fibrosis, and eosinophilia, accompanied by small subcutaneous fat lobules enclosed by thickened fibrous septa. The perivascular inflammatory infiltrate was formed by lymphocytes, plasma cells, and abundant eosinophils (Fig. 2a-c). These findings strongly suggested eosinophilic fasciitis (EF) rather than a drug-related reaction, considering the timeline between drug cessation and lesion onset, leading to a diagnosis of paraneoplastic eosinophilic fasciitis.

She underwent a 40-day prednisone treatment, beginning at 0.75 mg/kg/day and gradually tapered every 6 to 10 days. Within a month of therapy, her skin lesions completely resolved, accompanied by a notable decrease in eosinophil count (0.06/mm³) based on blood tests.

Three months later, eosinophilic fasciitis reappeared in the lower third of the left leg, confirmed by histopathological examination (Fig. 1b). During an oncology check-up, the patient was diagnosed with pelvic metastasis, leading to the commencement of doxorubicin and vinorelbine combined

therapy. Simultaneously, she received a 30-day course of oral prednisone at 0.75 mg/kg/day, gradually tapered, swiftly achieving remission of the skin lesions. At the 6-month follow-up, no further recurrences were noted, and there was no sign of disease progression.

Discussion and Conclusions

EF is a rare syndrome of unknown aetiology that causes scleroderma-like symptoms, characterized by skin and deeper perimuscular fascial plane induration associated with peripheral eosinophilia.

Various hematologic disorders, both benign and malignant, as well as drug-associated conditions, have been correlated with EF^{2,3}. Non-hematological malignancies such as melanoma, nonmelanoma skin cancer, breast, lung, thyroid, prostate, gastrointestinal, and ovarian cancer have been associated with eosinophilic fasciitis⁴.

According to Curth's postulates, based on a collection of clinical criteria used to assess the link between an underlying malignancy and dermatological conditions, EF may act as a paraneoplastic syndrome⁵. Frequently, paraneoplastic EF displays resistance to immunosuppressive treatments, including corticosteroids, while showing responsiveness to therapies targeting the underlying malignancy⁶.

Merely three instances of eosinophilic fasciitis in conjunction with breast cancer have been documented, where the onset of fasciitis closely preceded or followed the carcinoma diagnosis^{2,4}. Our case is in agreement with the literature.

In our case, the first episode of EF could be attributed to a paraneoplastic manifestation of recurrent underlying neoplastic pathology. The most intriguing aspect of our case is that the recurrence of EF coincided with the detection of metastatic secondary lesions, confirming the causal relationship even in cases of metastasis.

Our findings conclude that EF indicates a possible recurrence of neoplastic growth and should be considered a potential red flag. This underscores the importance of a dermatological evaluation in patients with EF and a concurrent history of neoplasia.

References

1. Long H, Zhang G, Wang L, Lu Q. Eosinophilic Skin Diseases: A Comprehensive Review. *Clin Rev Allergy Immunol*. 2016;50(2):189-213
2. Haddad H, Sundaram S, Magro C, Gergis U. Eosinophilic fasciitis as a paraneoplastic syndrome, a case report and review of the literature. *Hematol Oncol Stem Cell Ther*. 2014;7(2):90-92.

3. Maione V, Miccio L, Sala R, Zambelli C, Calzavara-Pinton P. Eosinophilic fasciitis induced by natalizumab in a patient affected by multiple sclerosis. *Indian J Dermatol Venereol Leprol.* 2021;87(1):146.
4. Watts RA, Merry P. Familial eosinophilic fasciitis and breast cancer. *Br J Rheumatol.* 1994;33(1):93-94.
5. Silva JA, Mesquita Kde C, Igreja AC, et al. Paraneoplastic cutaneous manifestations: concepts and updates. *An Bras Dermatol.* 2013;88(1):9-22.
6. Philpott H, Hissaria P, Warrren L, et al. Eosinophilic fasciitis as a paraneoplastic phenomenon associated with metastatic colorectal carcinoma. *Australas J Dermatol.* 2008;49(1):27-29.



Figure 1. a) Erythema, edema, and a "peau d'orange" texture on the left leg; b) recurrence of EF in the lower third of the left leg.




Figure 2. a) Skin biopsy of eosinophilic fasciitis (H&E, 5x); b) The perivascular inflammatory infiltrate includes lymphocytes, plasma cells and numerous eosinophils (H&E, 20x); c) Histopathology displaying hyperkeratotic epidermis, fibrosis and eosinophilia of the reticular dermis, while the subcutaneous fat lobules appear diminished in size and are encased by thickened fibrous septa (H&E, 10x).