



Dermatology Reports

<https://www.pagepress.org/journals/index.php/dr/index>

eISSN 2036-7406



Publisher's Disclaimer. E-publishing ahead of print is increasingly important for the rapid dissemination of science. **Dermatology Reports** is, therefore, E-publishing PDF files of an early version of manuscripts that undergone a regular peer review and have been accepted for publication, but have not been through the copyediting, typesetting, pagination and proofreading processes, which may lead to differences between this version and the final one.

The final version of the manuscript will then appear on a regular issue of the journal. E-publishing of this PDF file has been approved by the authors.

Please cite this article as: Veraldi S, Aromolo IF, Germiniasi FS, Nazzaro G. Erythrasmoid pityriasis versicolor: three case reports and review of the literature. Dermatol Rep 2024 [Epub Ahead of Print] doi: 10.4081/dr.2024.9976

 © the Author(s), 2024
Licensee PAGEPress, Italy

Submitted 24/02/2024 – Accepted 15/03/2024

Note: The publisher is not responsible for the content or functionality of any supporting information supplied by the authors. Any queries should be directed to the corresponding author for the article.

All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article or claim that may be made by its manufacturer is not guaranteed or endorsed by the publisher.

Erythrasmoid pityriasis versicolor: three case reports and review of the literature

Stefano Veraldi,¹ Italo Francesco Aromolo,² Francesca Sara Germiniasi,³ Gianluca Nazzaro²

¹Dermatological Centre, Milan; ²Department of Pathophysiology and Transplantation, Università degli Studi, IRCCS Foundation, Ca' Granda Ospedale Maggiore Policlinico, Milan; ³San Raffaele Hospital, Milan, Italy

Correspondence: Italo Francesco Aromolo, Dermatology Unit, Via Pace, 9, 20122 Milan, Italy.
Tel.: +39.3278752727.

E-mail: italo.aromolo@unimi.it

Key words: *Malassezia* sp.; pityriasis versicolor; erythrasmoid pityriasis versicolor; erythrasma.

Contributions: SV, IFA, wrote the manuscript; SV, IFA, FSG, data collection and interpretation; SV, GN, study concept, design, and supervision. All the authors approved the final version to be published.

Conflict of interest: the authors declare no potential conflict of interest.

Funding: none.

Ethical approval and consent to participate: the study was conducted in accordance with the ethical standards of the responsible committee on human experimentation (institutional and national), with the Helsinki Declaration of 1975, as revised in 2000, and with the Taipei Declaration.

Availability of data and material: data and materials are available from the corresponding author upon request.

Consent for publication: all enrolled patients and their parents provided written consent for the retrospective study of the collected data.

Abstract

Pityriasis versicolor is characterized clinically by slightly scaly, hyper- or hypopigmented macules usually located on the chest, shoulders, back and arms. Rare cases of atypical locations of pityriasis versicolor have been described, such as groins, penis and perineum. We present three cases of pityriasis versicolor located exclusively on the submammary folds, pubis and inguinal folds (the latter in an 11-year-old girl), respectively. In consideration of both location and clinical presentation of pityriasis versicolor in these three patients, we think that the diagnosis of “erythrasmoid” pityriasis versicolor is acceptable. Differential diagnosis with erythrasma must be taken into consideration. Mycological and bacteriological examinations are therefore necessary in order to make a correct diagnosis and specific therapy.

Introduction

Pityriasis versicolor (PV) (tinea versicolor) is a common, superficial skin infection caused by *Malassezia* sp.. The latter consists of several subspecies that belong to the normal cutaneous flora, in particular in areas which are rich in sebaceous glands. *Malassezia* sp. can convert from saprophyte to pathogenic yeast when some predisposing factors occur, such as hot-humid environment (PV is more frequent in Tropical and Subtropical countries), hyperhidrosis, organ transplants, therapies with antibiotics, corticosteroids and immunosuppressive drugs.¹

PV is characterized clinically by slightly scaly, hyper- or hypopigmented macules located on the chest, shoulders, back and arms.¹ Rare cases of atypical locations of PV have been described,²⁻²¹ such as groins,^{2,5-7,10,14,15,17, 18,21} penis^{2-4,8,9,11-14,16,19,20} and perineum.^{2,5,14} To our knowledge, no cases of PV involving exclusively the submammary folds have been reported and only one case of PV involving uniquely the pubis has been described.¹⁶

We present three cases of “erythrasmoid” PV located exclusively on the submammary folds, pubis and inguinal folds (the latter in an 11-year-old girl), respectively.

Case Report

Case 1

An 18-year-old Caucasian female was admitted because of a pigmentation located on the submammary folds. The patient stated that she was well and that was not in therapy with systemic drugs. Dermatitis appeared approximately one year earlier: it was unsuccessfully treated at other centres with topical corticosteroids.

Dermatological examination revealed two symmetrical macules located on the submammary folds: they were brownish in colour, with fine scaling and irregular, well defined borders (Figure 1). The patient complained of severe itching. No other lesions were observed elsewhere.

Laboratory examinations were within normal ranges. Wood's lamp examination revealed a yellow fluorescence. Microscopical examination with 10% potassium hydroxide showed several round spores and septate and short hyphae. Bacteriological examination was negative.

A diagnosis of PV located exclusively on the submammary folds was made. The patient was successfully treated with isoconazole cream (1 application/day for three weeks) and oral itraconazole (200 mg/day for ten days). Follow up (six months) was negative for recurrences.

Case 2

A 38-year-old Caucasian female was admitted because of a dermatitis located on the pubis. She was in good general health and she was not in therapy with systemic drugs. She declared that the dermatitis appeared approximately four months earlier and that it was not previously treated.

Dermatological examination showed a macule located on the pubis: it was orange-ochre in colour, with fine scaling and well defined borders (Figure 2). The patient complained of mild itching. No other lesions were observed elsewhere.

Laboratory examinations were within normal limits. Wood's lamp examination showed a yellowish fluorescence. Microscopical examination with 10% potassium hydroxide revealed several round yeast spores and septate and short hyphae. Bacteriological examination was negative.

A diagnosis of PV located exclusively on the pubis was made. The patient was successfully treated with a shampoo containing tioconazole and zinc pyrithione (1 cleaning/day for three weeks). No recurrence was observed during a six month follow up.

Case 3

An 11-year-old Caucasian girl was admitted because of a dermatitis located on the inguinal folds. The patient's parents stated that she was in good general health and that she was not in therapy with systemic drugs. The dermatitis appeared approximately three weeks earlier and it was previously, although unsuccessfully, treated with zinc oxide.

Dermatological examination showed several macules located on the inguinal folds and inner surfaces of the thighs: they were brownish in colour, with irregular borders (Figure 3). The patient complained of mild itching. No other lesions were observed elsewhere.

Laboratory examinations were within normal limits. Wood's lamp examination showed a yellowish fluorescence. Microscopical examination with 10% potassium hydroxide revealed several round yeast spores and septate and short hyphae. Bacteriological examination was negative.

A diagnosis of PV located exclusively on the inguinal folds and thighs was made. The patient was successfully treated with a shampoo containing tioconazole and zinc pyrithione (1 cleaning/day for three weeks). Follow up (four months) was negative.

Discussion and Conclusions

Submammary folds, pubis and inguinal folds are very rare locations of PV. As previously mentioned, only one case of PV located exclusively on the pubis was reported.¹⁶ In addition, no cases of PV involving uniquely the inguinal folds were described. In consideration of both location and clinical presentation of PV in these three patients, we think that the diagnosis of "erythrasmoid" PV can be acceptable.¹⁵ However, we must not forget the possibility that PV and erythrasma can coexist, as reported by some authors.¹⁸ As previously mentioned, the search for *Corynebacterium minutissimum* was negative in all patients. Mycological and bacteriological examinations are therefore necessary in order to make a correct diagnosis and specific therapy.

References

1. Veraldi S. Pityriasis versicolor. ISED Brescia (Italy) 1998;7-14.
2. Roberts SOB. Pityriasis versicolor: a clinical and mycological investigation. Br J Dermatol 1969;81:315-26.
3. Blumenthal HL. Tinea versicolor of penis. Arch Dermatol 1971;103:461-2.
4. Schaller KF. Mykosen des Intimbereichs. II. Dermatophytien, Pityriasis versicolor, Erythrasma und Trichomycosis palmellina. Mykosen 1971;14:447-9.
5. Rudolph RI, Holzwanger JM. Inverse tinea versicolor. Arch Dermatol 1975;111: 1213.
6. Kamalam A, Thambiah AS. A study of 3891 cases of mycoses in the tropics. Sabouraudia 1976;14:129-48.
7. Abdul Razack EM, Thambiah AS. A clinical study of pityriasis versicolor in Madras. Sabouraudia 1977;15:305-11.
8. Smith EL. Pityriasis versicolor of the penis. Br J Vener Dis 1978;54:441.
9. Nia AK, Smith EL. Pityriasis versicolor of the glans penis. Br J Vener Dis 1979;55: 230.
10. Burkhart CG, Dvorak N, Stockard H. An unusual case of tinea versicolor in an immunosuppressed patient. Cutis 1981;27:56-8.

11. Daneshvar SA, Hashimoto K. An unusual presentation of tinea versicolor in an immunosuppressed patient. *J Am Acad Dermatol* 1987;17(2 Pt 1):304-5.
12. Aljabre SHM, Sheikh YH. Penile involvement in pityriasis versicolor. *Trop Geogr Med* 1994;46:184-5.
13. Kaur I, Handa S, Kumar B. Tinea versicolor: involvement of unusual sites. *Int J Dermatol*. 1996; 35: 604-5.
14. Assaf RR, Weil ML. The superficial mycoses. *Dermatol Clin*. 1996; 14: 57-67.
15. Gorani A, Oriani A, Falconi Klein E, Veraldi S. Case report. Erythrasmoid pityriasis versicolor. *Mycoses* 2001;44:516-7.
16. Aljabre SHM. Intertriginous lesions in pityriasis versicolor. *J Eur Acad Dermatol Venereol* 2003;17:659-62.
17. Aste N, Pau M, Aste N. Pityriasis versicolor on the groin mimicking erythrasma. *Mycoses* 2004;47:249-51.
18. Karakatsanis G, Vakirlis E, Kastoridou C et al. Coexistence of pityriasis versicolor and erythrasma. *Mycoses* 2004;47:343-5.
19. Aridoğan IA, Ilkit M, Izol V et al. *Malassezia* and *Candida* colonisation on glans penis of circumcised men. *Mycoses* 2005;48:352-6.
20. Ryu HW, Cho JW, Lee KS. Pityriasis versicolor on penile shaft in a renal transplant recipient. *Ann Dermatol* 2012;24:345-7.
21. Varada S, Dabade T, Loo DS. Uncommon presentations of tinea versicolor. *Dermatol Pract Concept* 2013;4:93-6.



Figure 1. Pityriasis versicolor located exclusively on the submammary folds.



Figure 2. Pityriasis versicolor located exclusively on the pubis.



Figure 3. Pityriasis versicolor located on the inguinal folds in an 11-year old girl.