

Review on the consequences of using Improvac™ in modern pig production

Anna Katharina Albrecht

Institute for Animal Breeding and Husbandry, Christian Albrechts Universität, Kiel, Germany

Abstract

In Europe growing concerns regarding animal welfare issues in pig production have forced the pig industry to introduce alternative methods to conventional castration procedures. Besides the raising of entire males and castration in combination with analgesia and anaesthesia the vaccination against GnRF (Gonadotropin-releasing factor) seems to be the most promising long-term solution. Immunised male pigs (IM) show higher average daily weight gain than surgically castrated males (CM). Additionally the feed intake in IM is lower than in CM and feed conversion ratio is consequently better. Carcass weight, back fat depth and dressing percentage of IM pigs are intermediate between CM and entire males while meat quality seems not to be influenced by castration technique since CM and IM show comparable results. Steroid hormone concentrations in IM decline to very low levels (below detection line) after the second administration of the anti-GnRF vaccine and boar taint compounds are reliably metabolized. Pigs which received two injections of the anti-GnRF vaccine reduce their sexual and aggressive behaviour to levels of CM pigs which results in low incidents of injury and carcass damages. Surveys analyzing the consumers' attitude to vaccination against boar taint reveal that if profound information on the technique is provided, the acceptance of meat from vaccinated animals is even better than the acceptance of meat from pigs castrated under current farm conditions. Furthermore economic analyses reveal that immunisation against GnRF provides a potential for a return on investment since better feed efficiency compensates for the additional costs of drug and labour time.

Introduction

Over the last couple of years a large variety of alternative methods to surgical castration without anaesthesia has been presented. This development is mainly due to an increase in scientific knowledge on the phys-

iology of pain in young animals, an increase in the public awareness and concerns regarding the castration procedure and as a consequence an increase in the demand for a more animal-friendly castration technique and improved animal welfare. The conventional procedure, *i.e.* surgical castration without anaesthesia and analgesia within the first seven days of life, has been common practice in most European countries over the last centuries. The EU countries alone produce about 250 million slaughter pigs each year.¹ Castration, as a means of preventing boar taint and aggressive behaviour, is performed on 77% of male pigs.² Growing public concerns regarding animal welfare and changing legal requirements have forced governments and the pig industry to reconsider the traditional approach and to reinforce the effort to introduce alternative methods. The most recognised approach, besides raising of entire males and surgical castration with analgesia/anaesthesia, is the down-regulation of gonadal hormones by using vaccination against Gonadotropin-releasing factor (GnRF). The product focused on in this review is Improvac™ (Pfizer Animal Health) since immunisation against GnRF is the most favoured among the vaccination techniques and products.

In the interest of completeness it should be mentioned that other ways of down-regulating of gonadal hormone activity exist.

Castration via vaccination can either be directed against the pituitary hormone LH or the hypothalamic hormone GnRF. Both approaches usually use active immunisation, although passive immunisation is also possible, however, it has proven to be less effective.³ Falvo *et al.*⁴ compared the vaccination of boars with LH and GnRF vaccines and came to the conclusion that LH vaccination was less effective when compared with immunisation against GnRF. In the interest of completeness, it should be mentioned that vaccination against 5 α -Androstenone is also possible, but has also proven to be less effective.⁵ Over the last 30 years a large number of GnRF vaccines has been subjected to various studies as reviewed by Prunier *et al.*⁶ In modern production systems only a vaccine with manageable labour costs and good tolerance can prevail. Modern anti-GnRF vaccines use tolerable adjuvants and only two injections. With these vaccines two possible vaccination schedules exist, early and late vaccination. Studies conducted by Turkstra *et al.*⁷ and Zeng *et al.*⁸ used a vaccine which is administered relatively early during the pig's life. These vaccines hold the advantage of easier detection of successfully vaccinated animals at the slaughter line. The production advantages of entire male boars, however, were diminished in those pigs and the vaccinated animals showed a growth per-

Correspondence: Anna Katharina Albrecht, Institute for Animal Breeding and Husbandry, Christian Albrechts Universität, Olshausenstraße 40, 24098 Kiel, Germany.
Tel. +49.431.8802585 - Fax: +49.431.8802588.
E-mail: jkrieter@tierzucht.uni-kiel.de

Key words: vaccination, castration, boar taint.

Conflict of interests: the author declares no potential conflict of interests.

Received for publication: 5 October 2011.

Revision received: 30 November 2011.

Accepted for publication: 30 November 2011.

This work is licensed under a Creative Commons Attribution NonCommercial 3.0 License (CC BY-NC 3.0).

©Copyright A.K. Albrecht, 2013

Licensee PAGEPress srl, Italy

Veterinary Science Development 2013; 3:e1

doi:10.4081/bsd.2013.e1

formance and carcass characteristics comparable to those of barrows.

The most recognised late-vaccination technique is the immunisation against GnRF with two injections given at least 4 weeks apart with the second injection given four to six weeks prior to slaughter.

This review in particular focuses on the effects of immunisation against GnRF on growth performance, carcass characteristics and meat quality, blood testosterone concentrations and behaviour. Additionally, short insight is provided into the effects of using immunisation against GnRF on the major boar taint compounds androstenone and skatole, testes size, the consumer's acceptability of meat from vaccinated pigs and the economic implications of using immunisation against GnRF in modern pig production.

In the interests of completeness, it should be mentioned that castration via vaccination is not only used in male pigs but in a large variety of mammals (all species: Ferro *et al.*,⁹ Thompson *et al.*,¹⁰ cattle: Ribeiro *et al.*,¹¹ Bonneau and Enright,¹² ram lambs: Ülker *et al.*,^{13,14} goat bucks: Godfrey *et al.*¹⁵ In all of these species, the purposes of vaccination are more or less the same: the improvement of meat and carcass characteristics, a reduction in male aggressive behaviour, reduction in male-associated odour (esp. swine and goat), and in the case of pet species, the neutralisation of fertility.

Immunisation against GnRF has also been tried in females but only plays a tangential role and is not mentioned further.^{16,17}

Effects of using immunisation against Gonadotropin-releasing factor on growth performance

Most authors agree that treatment with anti-GnRF vaccines has no effect on growth performance before the second vaccination (V2) is administered and that these pigs [immunised male (IM) pigs] perform comparable to intact boars during the first part of the fattening period, *i.e.* before V2.^{18,21} Body weight and average daily weight gain (ADG) seem to be at comparable levels for IM pigs, boars and barrows [castrated males (CM) pigs] until V2,^{18,19,21,22} whereas the feed intake (FI) in CM pigs is higher when compared with the other groups.^{18,22} Hemonic *et al.*²² and Cronin *et al.*¹⁸ report lower average daily feed intake (ADFI) in IM pigs when compared to CM pigs during the first part of the fattening period. From literature it is well known that entire males show lower voluntary ADFI than CM pigs.¹⁸ Cronin *et al.*¹⁸ and Dunshea *et al.*¹⁹ argue that this lower ADFI results from the fact that entire males allocate more of their active time to social behaviour and spend less time feeding. Furthermore, Weiler *et al.*²³ found a negative correlation for voluntary ADFI and testosterone levels in the blood. IM pigs can be regarded as entire males until the second vaccination is given.^{19,20} Despite the lower voluntary ADFI in immunisation against GnRFTM-treated pigs, average daily gain (ADG) and body weight (BW) are comparable to those of surgical castrates until the time point of the V2.^{18,24} This results from the anabolic effect of the male steroid hormones, which are at comparable levels in IM males and intact boars before V2 (see below). After V2, however, IM pigs increase their voluntary ADFI to levels comparable to CM pigs or even higher.^{18,19,22,25-27} In contrast, Skrlep *et al.*²¹ and Pauly *et al.*²⁰ found that although ADFI increases after V2, IM pigs do not reach the levels of CM pigs but consume more feed than intact males of the same age. This increase in ADFI results from a change in the hormonal profile of IM pigs as well as a change in the pigs' behaviour.^{18,19} Cronin *et al.*¹⁸ found that intact males spend less time feeding than CM and vaccinated pigs after V2. Additionally, IM pigs after V2 alter their behaviour and spend less time on social, especially aggressive and sexual behaviour. As a consequence IM pigs reveal higher ADG. Dunshea *et al.*¹⁹ came to the conclusion that the higher ADG in IM pigs compared to CM pigs is a result of this increase in ADFI in combination with the more barrow-like behaviour rather than a better feed conversion ratio (FCR).

Effects of using immunisation against Gonadotropin-releasing factor on carcass characteristics and meat quality

Intact males are known to have lower carcass weight and dressing percentage than CM pigs.^{7,8,28} Most authors found intermediate values for IM pigs for both parameters.^{19,28} Gispert *et al.*²⁸ and Dunshea *et al.*¹⁹ explain these lower values in IM pigs with the higher gut fill and the removal of the testes. Another effect of vaccination against boar taint is the increase in lean meat percentage comparative to CM pigs.^{20,24,25,29,30} Along with this increase in lean meat percentage comes a reduction in intramuscular fat content and backfat thickness in IM pigs,²⁰ which show intermediate values when compared to entire males and CM pigs.^{1,20,28,29,31} The proportion of the ham, as analysed by Gispert *et al.*²⁸ and Pauly *et al.*,²⁰ and the proportion of the loin revealed higher values for entire males when compared to CM pigs,²⁰ with IM pigs in between. Meat quality seems not to be affected by vaccination.^{20,28,29} The compensatory growth and the reduction in intramuscular fat content were expected to be detrimental to meat quality parameters such as tenderness and juiciness. The study conducted by Pauly *et al.*,²⁰ however, found no evidence for reduced tenderness and drip loss in pork from IM pigs. Additionally, Dunshea *et al.*¹⁹ found that, independent of the period of time between V2 and slaughter, IM pigs that were slaughtered with 23 weeks of age had dressing percentages comparable to those of intact boars, whereas slaughter at an advanced age (26 weeks) revealed higher values for intact boars compared to IM pigs. Lealiifano *et al.*³² found that the timing of V2 had a great influence on many carcass characteristics. Those pigs which received the second immunisation four to six weeks prior to slaughter showed carcass values similar to CM pigs, whereas pigs given a late vaccination, *i.e.* two weeks before slaughter maintained many of the performance advantages of intact boars. Rikard-Bell *et al.*²⁷ state that the increase in ADFI in IM pigs after V2 results in a great deal of that additional energy being converted into fat rather than muscle growth. They further found that combination of immunisation against GnRF-treatment and ractopamine had additive effects not only on growth performance but that carcass composition was positively influenced, since ractopamine is a stimulator of adipose tissue fat mobilisation. IM pigs which received vaccination as well as ractopamine supplementation revealed an increase in carcass weight and lean meat percentage, whereas half carcass fat mass and backfat thickness had decreased. Similar effects were reported by Oliver *et al.*,²⁶ who

analysed the effects of vaccination against boar taint in combination with porcine somatotropin (pST). Porcine somatotropin is a peptide hormone used to alter the partitioning of energy in feed away from fat and towards muscle growth. Both compounds (ractopamine and pST) are not approved as feed additives in the EU.

Effects of using immunisation against Gonadotropin-releasing factor on testosterone levels in the blood

Testosterone levels in the blood seem to follow a similar pattern to androstenone concentrations in the adipose tissue.³² Until V2, testosterone levels in the blood of IM pigs are comparable to those of entire male pigs,^{19,30,33} which show increasing concentrations with age. CM pigs, on the contrary, show testosterone concentrations below the detection limit.³³ IM pigs reveal a significant decrease in the blood testosterone concentration after V2.^{19,33} At slaughter, IM pigs regularly display testosterone concentrations comparable with CM, *i.e.* at very low levels or below detection limit.^{19,22,30} Only few studies have focused on the long-term effects of immunisation against GnRF. Zamaratskaia *et al.*³³ found that the effects of immunisation against GnRF on hormonal profile lasted until at least 22 weeks after V2, at which time testosterone levels in IM pigs were still at lower levels than in entire boars.

Effects of using immunisation against Gonadotropin-releasing factor on behaviour and animal welfare

The results on the behavioural consequences of vaccination against boar taint are very consistent, although only few studies have so far examined the effects of immunisation against GnRF on behaviour, especially aggressive and sexual behaviour. Most authors agree, that entire males and IM pigs spend more time on social and active behaviour (as indicated by the number of standing, walking or eating pigs) than CM pigs in the period before V2.^{18,33-35} After V2, IM pigs alter their behaviour significantly and perform social and active behaviour at comparable levels with CM pigs or females and differ significantly from intact boars.^{34,18,25,36} The reduction in active behaviour is also associated with a reduction in aggressive, mounting and sexual behaviour.

Whereas IM pigs before V2 perform aggressive and mounting behaviour at levels comparable with intact males and at much higher levels than CM pigs, the second immunisation against GnRF leads to a significant decrease in these behavioural traits as soon as one week after V2.^{18,25,34,36} Additionally, a study conducted by Zamaratskaia *et al.*,³⁰ examining the long-term effects of vaccination, revealed that these changes last up to 21 weeks after V2.

This reduction in aggressive and mounting behaviour further results in fewer skin lesions in immunised pigs in comparison to intact boars at slaughter.^{25,36} Rydhmer *et al.*³⁶ state that most of the skin lesions found in intact boars at slaughter result from mounting rather than fighting activities.

Aggressive and sexual behaviour are important indicators of animal welfare since high levels of aggression and mounting behaviour impose stress, fear and injury not only on the receiver but on all pigs in the pen.³⁷ Animal welfare consequences of surgical castration and its alternatives have been reviewed further by Prunier *et al.*⁶ and Borrell *et al.*³⁸ Both authors come to the conclusion that vaccination against boar taint offers a good alternative to surgical castration since not only the pain and discomfort associated with the procedure are avoided but fighting behaviour is reduced after V2.

Effects of using immunisation against Gonadotropin-releasing factor on the boar taint compounds androstenone and skatole, testes size, consumer's acceptability and economic implications

The effects of vaccination against boar taint have been evaluated by many studies since boar taint has been the major reason for castration in the past. Extensive studies as reviewed by Xue *et al.*³⁹ and Stefan Guizot,⁴⁰ have identified androstenone and skatole as major contributors to boar taint. Numerous studies have proven that immunisation against GnRF is very effective in reducing boar taint,^{1,18,19,24,30,41} since androstenone and skatole are reliably metabolized in the period after V2.^{21,22,24,32} Lealiifano *et al.*³⁰ further found that even pigs slaughtered only two weeks after V2 show androstenone and skatole levels comparable to barrows and well below threshold limits. Along with the reduction in the concentration of male steroid hormones comes a reduction in the size of the reproductive organs.^{21,22,28,42,43} Some authors suggested using the size of the testicles as an indicator of successful vaccination.¹⁹ However, since not

only the genetic background and the age at slaughter influence testicle size but also the time between V2 and slaughter a reliable detection of tainted pork by testes size alone seems impossible.^{1,20,31,32,44}

The major reason for the close examination of boar taint and the importance of detecting tainted meat at the slaughter line are the potential of such tainted meat to cause taste and smell aberrations in heated pork which most consumers strongly object to.^{45,46} Surveys conducted in order to gain knowledge on the acceptance of tainted meat on the one hand, and meat from IM males, on the other, have been carried out in many countries.⁴⁵⁻⁵¹ The acceptance of tainted meat is very poor in most European countries although differences exist. These differences in the consumers' acceptability can be due to different cooking and evaluation methods as well as to the consumers' origins, ages, sex or androstenone sensitivities.^{45,52-54} The acceptance of meat from IM pigs differ among the studies. Font I Furnols *et al.*⁴⁵ found that there was no significant difference in the evaluation of meat from immunised pigs, surgically castrated pigs and females and came to the conclusion that the products of immunised males were indistinguishable from pork from barrows or females. The study conducted by Huber-Eicher and Spring revealed that most consumers have no clear association with the term *immunocastration*,⁴⁶ but that meat from IM pigs would be much less accepted than meat from pigs surgically castrated under anaesthesia. Hofer and Kupper conducted a survey on more informed consumers.⁵⁰ The participants were given information on the actual situation of castration and the alternatives. The results of degustation were in line with the findings of Font I Furnols *et al.*⁴⁵ More information on vaccination against boar taint, however, seemed to have had positive influence on the consumers' acceptability of meat from immunisation against GnRF-treated pigs, since the majority agreed on vaccination as a feasible alternative to current practice.

For any alternative method to surgical castration it is necessary to evaluate its economic effects, since only methods can prevail which have few financial disadvantages for the stakeholders. Deen *et al.*⁵⁵ come to the conclusion that immunisation against GnRF offers potential for a return on investment. However, the financial effects must always be contextualised within the constraints of each production system since the production implications will vary in different systems. De Roest *et al.*⁵⁶ came to the conclusion, looking at the EU countries only, that the better feed efficiency of vaccinated pigs can compensate for the costs of vaccination. A lot, however, depends on the costs of the vaccine and the consumers' acceptance of the procedure.

Conclusions

In summary, most authors agree on vaccination against boar taint as a feasible alternative to surgical castration, since its effectiveness in preventing boar taint has been reliably proven in many studies. In addition, vaccination avoids surgical procedures, is effective in decreasing the occurrence of fighting and mounting behaviour and may improve the feed conversion ratio and lean meat percentage without having adverse effects on meat quality parameters.

References

1. Fuchs T, Nathues H, Koehrmann A, et al. A comparison of the carcass characteristics of pigs immunized with a gonadotrophin-releasing factor (GnRF) vaccine against boar taint with physically castrated pigs. *Meat Science* 2009;83:702-5.
2. Fredriksen B, Hexeberg C. The effect of removing animals for slaughter on the behaviour of the remaining male and female pigs in the pen. *Res Vet Sci* 2009;86:368-70.
3. Van der Lende T, Kruijt L, Tieman M. Can passive immunisation with anti-GnRF monoclonal antibodies, injected a few weeks before slaughter, prevent boar taint? In: Bonneau M, ed. *Measurement and prevention of boar taint*. Paris: INRA Editions; 1993. pp 201-206.
4. Falvo RE, Chandrashekar V, Arthur RD, Kenstler, et al. Effect of active immunisation against LHRH or LH in boars: reproductive consequences and performance traits. *J Anim Sci* 1986;63:986-94.
5. Williamson ED, Patterson RLS. A selective immunisation procedure against 5 α -androstenone in boars. *Anim Product* 1982;35:353-60.
6. Prunier A, Bonneau M, von Borell EH, et al. A review of the welfare consequences of surgical castration in piglets and the evaluation of non-surgical methods. *Anim Welfare* 2006;15:277-89.
7. Turkstra JA, van Diepen JTM. Performance of male pigs immunised against GnRF is related to the time of onset of biological response. *J Anim Sci* 2002;80:2953-9.
8. Zeng XY, Turkstra JA, Jongboed AW, et al. Performance and hormone levels of immunocastrated, surgically castrated and intact male pigs fed ad libitum high- and low-energy diets. *Livest Prod Sci* 2002;77:1-11.
9. Ferro VA, Khan MAH, McAdams D, et al. Efficacy of an anti-fertility vaccine based on mammalian gonadotrophin releasing

- hormone (GnRF-I)- a histological comparison in male animals. *Vet Immunol Immunopathol* 2004;101:73-86.
10. Thompson Jr DL. Immunisation against GnRF in male species (comparative aspects). *Anim Reproduct Sci*, 2000;60: 459-69.
 11. de Ribeiro EL, Hernandez JA, Zanella EL, et al. Growth and carcass characteristics of pasture fed LHRH immunocastrated, castrated and intact *Bos indicus* bulls. *Meat Sci* 2004;68:285-90.
 12. Bonneau M, Enright WJ. Immunocastration in cattle and pigs. *Livest Prod Sci* 1995;42:193-200.
 13. Ülker H, Kanter M, Gökdal Ö, et al. Testicular development, ultrasonographic and histological appearance of the testis in ram lambs immunized against recombinant LHRH fusion proteins. *Anim Reproducti Sci* 2002;86:205-19.
 14. Ülker H, Gökdal Ö, Temur C, et al. The effect of immunisation against LHRH on body growth and carcass characteristics in Karaka ram lambs. *Small Ruminant Res* 2002;45:273-8.
 15. Godfrey SI, Walkden-Brown SW, Martin GB, Speijers EJ. of goat bucks against GnRF to prevent seasonal reproductive and agonistic behaviour. *Anim Reproduct Sci* 1996;44:41-54.
 16. Robbins SC, Jelinski MD, Stotish RL. Assessment of the immunological and biological efficacy of two different doses of a recombinant GnRF vaccine in domestic male and female cats (*felis catus*) *J Reproduct Immunol* 2004;64:107-19.
 17. Zeng XY, Turkstra JA, Tsigos A, et al. Effects of active immunisation against GnRF on serum LH, inhibin A., sexual development and growth rate in Chinese female pigs. *Theriogenology* 2002;58:1315-26.
 18. Cronin GM, Dunshea FR, Butler KL, et al. The effects of immuno- and surgical- castration on the behaviour and consequently growth of group-housed male finisher pigs. *Appl Anim Behav Sci* 2003;81:111-26.
 19. Dunshea FR, Colantoni C, Howard K, et al. Vaccination of boars with a GnRF vaccine (immunisation against GnRF) eliminates boar taint and increases growth performance. *J Anim Sci* 2001;79:2524-35.
 20. Pauly C, Spring P, et al. Growth performance, carcass characteristics and meat quality of group-penned surgically castrated, immunocastrated (immunisation against GnRF (R)) and entire male pigs and individually penned entire male pigs. *Animal* 2009;3:1057-66.
 21. Skrlep MB, Segula B, Prevotnik M, et al. Effect of immunocastration (immunisation against GnRF) in fattening pigs i: growth performance, reproductive organs and malodorous compounds. *Slo Vet Res* 2010;47:57-64.
 22. Hemonic A, Courboulay V. Evaluation of the safety, efficacy and production benefits of vaccination against boar taint in male pigs raised under commercial field conditions in France. *Revue De Medicine Veterinaire* 2009;160:383-93.
 23. Weiler U, Claus R, Dehnhard M, Hofacker S. Influence of the photoperiod and a light reverse program on metabolically active hormones and food intake in domestic pigs compared with a wild boar. *Canad J Anim Sci* 1996;76:531-9.
 24. Jaros P, Burgi E. Effect of active immunisation against GnRF on androstenone concentration, growth performance and carcass quality in intact male pigs. *Livest Product Sci* 2005;92:31-8.
 25. Fabrega E, Velarde A, Cros J, et al. Effect of vaccination against gonadotrophin-releasing hormone, using immunisation against GnRF, on growth performance, body composition, behaviour and acute phase proteins. *Livest Product Sci* 2010;132:53-9.
 26. Oliver WT, McCauley I, Harrell RJ, et al. A gonadotropin-releasing factor vaccine (immunisation against GnRFTM) and porcine somatotropin have synergistic and additive effects on growth performance in group-housed boars and gilts. *J Anim Sci* 2003;81:1959-66.
 27. Rikard-Bell C, Curtis MA, van Barneveld RJ, et al. Ractopamine hydrochloride improves growth performance and carcass composition in immunocastrated boars, entire boars, and gilts. *J Anim Sci* 2009;87: 3536-43.
 28. Gispert M, Oliver MA, Velarde A, et al. Carcass and meat quality characteristics of immunocastrated male, surgically castrated male, entire male and female pigs. *Meat Sci* 2010;85:664-70.
 29. D'Souza DN, Mullan BP. The effect of genotype and castration method on the eating quality characteristics of pork from male pigs. *J Anim Sci* 2003;77:67-72.
 30. Zamaratskaia G, Andersson HK, Chen G, et al. Effect of a gonadotropin-releasing hormone vaccine (immunisation against GnRFTM) on steroid hormones, boar taint compounds and performance in entire male pigs. *Reprod Domest Anim* 2008; 43:351-9.
 31. Schmoll F, Kauffhold J, Pfuetzner A, et al. Growth performance and carcass traits of boars raised in Germany and either surgically castrated or vaccinated against gonadotropin-releasing hormone. *J Swine Health Product* 2009;17:250-5.
 32. Lealiifano AK, Pluske JR, Nicholls RR, et al. Altering the timing of an immunocastration vaccine to optimise pig performance. In: van Barneveld RJ, ed. *Manipulating pig production XII*. Werribee: Australasian Pig Science Association; 2009. p 184.
 33. Zamaratskaia G, Rydhmer L, Andersson HK, et al. Long-term effect of vaccination against gonadotropin-releasing hormone, using immunisation against GnRF, on hormonal profile and behaviour of male pigs. *Anim Reproduct Sci* 2007;108:37-48.
 34. Baumgartner J. The behaviour of male fattening pigs following either surgical castration or immunisation with a GnRF vaccine. *Appl Anim Behav Sci* 2010;124:28-34.
 35. Velarde A, Gispert M, Oliver MA, et al. The effect of immunocastration on the behaviour of pigs. *Proceedings of the 41st International Congress of the International Society for Applied Ethology*. 2007, 8-12 August. Merida, Mexico.
 36. Rydhmer L, Lundström K, Andersson K. Immunocastration reduces aggressive and sexual behaviour in male pigs. *Animal* 2010;6:965-72.
 37. Rydhmer L, Zamaratskaia G, Andersson HK, et al. Aggressive and sexual behaviour of growing and finishing pigs reared in groups, without castration. *Acta Agric Scand Section A: Anim Sci* 2006;56:109-19.
 38. von Borrell E, Baumgartner J, Giersing M, et al. Animal welfare implications of surgical castration and its alternatives in pigs. *Cambridge: Cambridge University Press*; 2009. pp 1488-1496.
 39. Xue JL, Dial GD, Pettigrew JE. Performance, carcass, and meat quality advantages of boars over barrows: a literature review. *J Swine Health Production* 1997;5:21-8.
 40. Guizot S. A review of the causes and control of boar taint. 2009. Available from: <http://www.sapork.biz/a-review-of-the-causes-and-control-of-boar-taint/>
 41. McCauley I, Watt M, Suster D, et al. A GnRF vaccine (immunisation against GnRF (R)) and porcine somatotropin (Reporcin (R)) have synergistic effects upon growth performance in both boars and gilts. *Aust J Agricult Res* 2003;54:11-20.
 42. Einarsson S, Brunius C, Wallgre M, et al. Effects of early vaccination with Improvac (R) on the development and function of reproductive organs of male pigs. *Anim Reprod Sci* 2011;127:50-5.
 43. Fuchs T, Thun R, Parvizi N, et al. Effect of a gonadotropin-releasing factor vaccine on follicle-stimulating hormone und luteinizing hormone concentrations and on the development of testicles and the expression of boar taint in male pigs. *Theriogenology* 2009;72:672-80.
 44. Prunier A. Genital-tract development and 5-alpha-androstenone concentration pattern during growth in male and female

- pigs of european and chinese breeds. *Annales De Zootechnie* 1987;3:49-56.
45. Font i Furnols M, Gispert M, Guerrero L, et al. Consumers' sensory acceptability of pork from immunocastrated male pigs. *Meat Sci* 2008;80:1013-8.
 46. Huber-Eicher B, Spring P. Attitudes of Swiss consumers towards meat from entire or immunocastrated boars: a representative survey. *Res Vet Sci* 2008;85:625-7.
 47. Allison J. Immunisation against GnRF: Consumer acceptance. Proceedings Pfizer Symposium at 20th Into Pig Vet Soc Cong, Durban, South Africa 2009.
 48. Bonneau M, Walstra P, Claudi-Magnussen C, et al. An international study on the importance of androstenone and skatole for boar taint: IV. Simulation studies on consumer dissatisfaction with entire male pork and the effect of sorting out carcasses on the slaughter line, main conclusions and recommendations. *Meat Sci* 2000;54: 285-95.
 49. Giffin B. Consumer acceptance of the use of vaccination to control boar taint. In Proceedings 20th Int Pig Vet Soc Cong, Durban, South Africa 2008.
 50. Hofer S, Kupper T. Survey on the acceptance of the vaccination against boar taint. 2008 Available from: http://www.shl.bfh.ch/fileadmin/docs/Forschung/KompetenzenTeams/NutztierUmweltInteraktion/ProSchwein/Synthesebericht/B14_Umfrage_zur_Akzeptanz_der_Impfung_20080331.pdf
 51. Lagerkvist CJ, Carlsson F, Viske D. Swedish consumer preferences for animal welfare and biotech: a choice experiment. *AgBioForum* 2006;9:51-8.
 52. Agerhem H, Tornberg E. A comparison of the off-flavour of shoulder butts from entire male and female pigs using expert and consumer panels. Proceedings 40TH international congress of meat science and technology. 1994 The Hague, The Netherlands.
 53. Wood JD, Nute GR, Fursey GAJ, Cuthbertson A. The effect of cooking conditions on the eating quality of pork. *Meat Sci* 1195;40:127-35.
 54. Weiler U, Font I Furnols M, Fischer K, et al. Influence of differences in sensitivity of Spanish and German consumers to perceive androstenone on the acceptance of boar meat differing in skatol and androstenone concentrations. *Meat Sci* 2000;54:297-304.
 55. Deen J, O'Connor J, Sorensen S, Baker T. An economic model to assess costs of immunisation against GnRF to the swine producer for control of boar taint. In Proceedings of the 20th International Pig Veterinary Society (IPVS) Congress. 22-26 June 2008, Durban, South Africa.
 56. de Roest K, Montanari C, Fowler T, Baltussen W. Resource efficiency and economic implications of alternatives to surgical castration without anaesthesia. *Animal* 2009;3:1522-31.

Non-commercial use only